

1. The following are unrelated: (15 pts)

(a) Rewrite each of the following without the absolute value symbol:

i.  $|j^2 - 6j|$

**Solution:**

Since  $j > 3$ ,  $j^2 > 6j$ ,  $j^2 - 6j > 0$  so  $|j^2 - 6j| = \boxed{j^2 - 6j}$

ii.  $|j^{\sqrt{2}} - 2j|$

**Solution:**

Since  $j^{\sqrt{2}} < 2$ ,  $j^{\sqrt{2}} - 2 < 0$  hence  $|j^{\sqrt{2}} - 2j| = \boxed{j^{\sqrt{2}} - 2}$  or  $2 - j^{\sqrt{2}}$

(d) Add and simplify:  $\frac{2}{9} + \frac{5}{12} + 7^0$

**Solution:**

$$\frac{2}{9} + \frac{5}{12} + 7^0 = \frac{14}{9} + \frac{5}{12} + 1 \quad (4)$$

$$= \frac{56}{36} + \frac{15}{36} + \frac{36}{36} \quad (5)$$

$$= \frac{107}{36} \quad (6)$$

(e) Simplify:  $\frac{j}{2j} - \frac{7}{4j} + \frac{3j + j2j}{4j}$

**Solution:**

$$\frac{j}{2j} - \frac{7}{4j} + \frac{3j + j2j}{4j} = \frac{10 + 2}{8} \quad (7)$$

$$= \frac{12}{8} \quad (8)$$

$$= \frac{3}{2} \quad (9)$$

2. The following are unrelated. Leave your answers without negative exponents. (20 pts)

(a)  $(5b^3)^2 7a^3 a^6$

**Solution:**

$$(5b^3)^2 7a^3 a^6 = (5^2)(b^3)^2 7a^3 \quad (10)$$

$$= 25b^6 7a^3 \quad (11)$$

$$= 175b^6 a^3 \quad (12)$$

(b) Simplify:  $\frac{\sqrt[3]{32x^2}}{\sqrt[3]{2^4 \cdot 16}}$

**Solution:**

$$\frac{\sqrt[3]{32x^2}}{\sqrt[3]{2^4 \cdot 16}} = \frac{jxj \sqrt[3]{2 \cdot 16}}{\sqrt[3]{2^4 \cdot 4}} \quad (13)$$

$$= \frac{jxj4 \sqrt[3]{2}}{\sqrt[3]{2^6 \cdot 2}} \quad (14)$$

$$= 2jxj \quad (15)$$

(c) Simplify:  $\frac{2(x^2 y^3)^3}{8x^3 y^{1-3}}$

**Solution:**

$$\frac{2(x^2 y^3)^3}{8x^3 y^{1-3}} = \frac{2x^6 y^9}{8x^3 y^{-2}} \quad (16)$$

$$= \frac{x^3 y^{27}}{4y^{\frac{1}{3}}} \quad (17)$$

$$= \frac{y^{\frac{28}{3}}}{4x^3} \quad (18)$$

(d) Multiply to rewrite as a polynomial:  $\sqrt{x-1} + 3 \sqrt{x-1} - 3$

**Solution:**

$$\sqrt{x-1} + 3 \sqrt{x-1} - 3 = \sqrt{x-1}^2 - 3^2 \quad (19)$$

$$= x - 1 - 9 \quad (20)$$

$$= \boxed{x - 10} \quad (21)$$

3. The following are unrelated: (12 pts)

(a) Find the domain of the expression (give your answer in interval notation):  $\sqrt{\frac{x^2 - 9}{x(x-3)}}$

**Solution:**  $x - 3 = 0$  results in division by zero so  $x = 3$  must be excluded from the domain. As well  $x > 0$  since the square root of a negative number does not exist. So the answer is:  $(0; 3) \cup (3; \infty)$

(b) Combine into a single fraction:  $\frac{1}{x^2 - x - 2} - \frac{3}{x + 1}$

**Solution:**

$$\frac{1}{x^2 - x - 2} - \frac{3}{x + 1} = \frac{1}{(x - 2)(x + 1)} - \frac{3}{x + 1} \quad (22)$$

$$= \frac{1 - 3(x - 2)}{(x - 2)(x + 1)} \quad (23)$$

$$= \boxed{\frac{7 - 3x}{(x - 2)(x + 1)}} \quad (24)$$

(c) Evaluate the expression:  $\frac{1}{2}x^2 - x + 1$  when  $x = 4$

**Solution:**  $\frac{1}{2}(4)^2 - 4 + 1 = 8 - 4 + 1 = 5$

4. The following are unrelated: (12 pts)

(a) Simplify:  $\frac{3x^2 + 12}{x^2 - 5x}$

(b) Simplify the compound fraction:  $\frac{\frac{3}{x^2} - \frac{1}{x}}{\frac{9}{x^2} - 1}$

**Solution:**

$$\frac{\frac{3}{x^2} - \frac{1}{x}}{\frac{9}{x^2} - 1} = \frac{\frac{3-x}{x^2}}{\frac{9-x^2}{x^2}} \quad (27)$$

$$= \frac{3-x}{9-x^2} \quad (28)$$

$$= \frac{3-x}{(3-x)(3+x)} \quad (29)$$

$$= \boxed{\frac{1}{3+x}} \quad (30)$$

(c) Factor by grouping:  $9x^3 - 18x^2 - 4x + 8$

**Solution:**

$$9x^3 - 18x^2 - 4x + 8 = 9x^2(x-2) - 4(x-2) \quad (31)$$

$$= (x-2)(9x^2 - 4) \quad (32)$$

$$= (x-2)(3x)^2 - 2^2 \quad (33)$$

$$= \boxed{(x-2)(3x+2)(3x-2)} \quad (34)$$

5. Is  $x = -2$  a solution of the inequality  $x^3 - 2x < 2$

7. Solve each of the following equations. If there are no solutions write NO SOLUTIONS: (10 pts)

(a)  $\sqrt[3]{8y+2} = y - 4$

**Solution:**

$$\sqrt[3]{8y+2} = y - 4 \tag{40}$$

$$\sqrt[3]{8y+2} = y - 4 \tag{41}$$

$$8y+2 = (y-4)^3 \tag{42}$$

$$y^3 - 11y + 28 = 0 \tag{43}$$

$$(y-7)(y-4) = 0 \tag{44}$$

$$y = 4; 7 \tag{45}$$

Plugging into the original equation, we find  $y = 4$  to be extraneous. Hence  $y = 7$

(b) Solve for  $h$ :  $P = A + hdg$

**Solution:**

$$P = A + hdg \tag{46}$$

$$hdg = P - A \tag{47}$$

$$h = \frac{P - A}{dg} \tag{48}$$

8. Solve the following inequalities. Justify your answers by using a number line or sign chart if needed. Answers without full justification will not receive full credit. Express all answers in interval notation. (8 pts)

(a)  $3x + 1 < 6$

**Solution:**

$$3x + 1 < 6 \tag{49}$$

$$3x < 5 \tag{50}$$

$$x < \frac{5}{3} \tag{51}$$

Hence the interval of solution is  $\left(\frac{5}{3}; 7\right)$

$$(b) x^3 - 3x^2 = 0$$


**Solution:**

We start by factoring the left hand side, and then make use of a number line/sign chart to choose the relevant interval of solution

$$x^3 - 3x^2 = 0 \tag{52}$$

$$x^2(x - 3) = 0 \tag{53}$$

Setting the left side equal to zero we get two values that make the left side zero:  $x = 0$  and  $x = 3$ . Placing these on a number line and picking test values we obtain the following chart



Notice that  $x = 0$  is a solution. Hence the solution is  $0 \leq x < 3$ .

9. Find all the solutions to the following equation, including the complex solutions (Hint: factoring will be important)  $z^3 = 1$ . (5 pts)

**Solution:**

$$z^3 - 1 = 0 \tag{54}$$

$$(z - 1)(z^2 + z + 1) = 0 \tag{55}$$

From which we conclude that  $z = 1$  or

$$z = \frac{1 \pm \sqrt{(1)^2 - 4(1)(1)}}{2(1)} \tag{56}$$

$$= \frac{1 \pm \sqrt{-3}}{2} \tag{57}$$

$$= \frac{1 \pm \sqrt{3}i}{2} \tag{58}$$