

Department of Applied Mathematics

Final Exam

August 2015

1. Root

2. Quadrature

1. Root

Formulate Newton's method for solving the nonlinear  $2 \times 2$  system of equations

$$\begin{cases} f(x, y) = 0 \\ g(x, y) = 0 \end{cases}$$

In the same style as how one proves quadratic convergence in the scalar case for  $f(x) = 0$  show quadratic convergence (assuming sufficient smoothness of  $f, g$ , root being simple, etc.) in the  $2 \times 2$  case. Assuming the root  $x = \alpha, y = \beta$  to be of multiplicity one, define  $\epsilon_n = x_n - \alpha, \eta_n = y_n - \beta$ , and show that both  $\epsilon_{n+1}$  and  $\eta_{n+1}$  are of size  $O(\epsilon_n^2, \eta_n^2)$

2. Quadrature

Consider the quadrature formula

$$I_{\text{quad}} = \sum_{i=0}^n \alpha_i f(x_i) \quad x_i \in [-1, 1] \quad (1)$$

for the integral

$$I = \int_{-1}^1 f(x) w(x) dx,$$

where  $w(x)$

3. Lab 1


Assuming that  $\varphi_n, n=0, \dots$  form a set of orthogonal polynomials of degrees  $n$  over some interval  $[a, b]$  with weight function  $w(x) > 0$ , show that they obey a three-term recursion relation of the form

$$\varphi_{n+1}(x) = (a_n x + b_n) \varphi_n(x) + c_n \varphi_{n-1}(x) \quad n=1, 2, \dots$$

where the coefficients  $a_n, b_n, c_n$  do not depend on  $x$ .

4. Lab 2

Let  $A \in \mathbb{C}^{n \times n}$  be a symmetric complex valued matrix,  $A = A^T$ . It is possible to show that one can find vectors  $u$  and nonnegative numbers  $\mu$  solving the so-called



6. Non-PDE

Consider the Poisson's equation

$$(\partial_{xx} + \partial_{yy}) u = f(x, y) \quad \text{in } B = \{x^2 + y^2 < 1\}$$

with the Dirichlet boundary condition

$$u|_{(x,y) \in \partial B} = 0$$

Set  $f$  to be

$$f(x, y) = 4\pi^2 \cos(\pi x) \cos(\pi y) - 4\pi^2 \sin(\pi x) \sin(\pi y)$$

yielding the solution

$$u(x, y) = \cos(\pi x) \cos(\pi y) - \sin(\pi x) \sin(\pi y)$$

At a first glance it may appear that seeking a solution as a sine series,

$$u(x, y) = \sum_{m, n=1}^{\infty} u_{mn} \sin(m\pi x) \sin(n\pi y)$$

should be an efficient approach. However, it turns out that the sine series converges rather slowly.

- (a) Can you figure out why the convergence of the sine series is fairly slow?
- (b) What are other bases one can use to achieve high accuracy? Suggest a basis that would be more efficient in this case.
- (c) Sketch a numerical scheme to compute the solution with high accuracy.