

O . . . **a** . . . **a** . . . **a** . . . **a** . . .

Kathleen Finlinson,¹ Woodrow L. Shew², Daniel B. Larremore^{3,4} and Juan G. Restrepo^{1,*}

¹*Department of Applied Mathematics, University of Colorado Boulder, Boulder, Colorado 80309, USA*

²*Department of Physics, University of Arkansas, Fayetteville, Arkansas 72701, USA*

³*Department of Computer Science, University of Colorado at Boulder, Boulder, Colorado 80309, USA*

⁴*BioFrontiers Institute, University of Colorado Boulder, Boulder, Colorado 80303, USA*



to a target state but in minimizing the error when trying to maintain a macroscopic variable (e.g. ϵ).

In addition, the error computed theoretically from the activity S found by solving Eq. (8) (symbols) agrees qualitatively with the one obtained numerically from simulations of the full system (solid line), with quantitative agreement except for large values of S . Networks with a homogeneous degree distribution and $\langle k \rangle = 1$ have the largest controllable range.

values and the range of control strengths that yield stable control. While heterogeneity can be beneficial for robustness to random node failures [34], our results suggest that a more homogeneous degree distribution might be preferable for situations where control of a large range of macroscopic network activity levels is important.

A common critique of the hypothesis that the cerebral cortex may operate near criticality is that critical dynamics are

too noisy, as reflected in the large fluctuations in Fig. 1(a). For many aspects of brain function it is easy to imagine that these large fluctuations would cause trouble. However, our primary result here is that the noisy dynamics of criticality are, in fact, easy to control. This suggests that a brain might be able to take advantage of the other functional benefits of criticality while controlling its own noise to remain at a manageable level.

-
- [1] S. Yu, T. L. Ribeiro, C. Meisel, S. Chou, A. Mitz, R. Saunders, and D. Plenz, Maintained avalanche dynamics during task-induced changes of neuronal activity in nonhuman primates, *eLife* **6**, e27119 (2017).
 - [2] G. Scott, E. D. Fagerholm, H. Mutoh, R. Leech, D. J. Sharp, W. L. Shew, and T. Knöpfel, Voltage imaging of waking mouse cortex reveals emergence of critical neuronal dynamics, *J. Neurosci.* **34**, 16611 (2014).
 - [3] W. L. Shew, W. P. Clawson, J. Pobst, Y. Karimipannah, N. C. Wright, and R. Wessel, Adaptation to sensory input tunes visual cortex to criticality, *Nat. Phys.* **11**, 659 (2015).
 - [4] T. Bellay, A. Klaus, S. Seshadri, and D. Plenz, Irregular spiking of pyramidal neurons organizes as scale-invariant neuronal avalanches in the awake state, *eLife* **4**, e07224 (2015).
 - [5] Z. Ma, G. G. Turrigiano, R. Wessel, and K. B. Hengen, Cortical circuit dynamics are homeostatically tuned to criticality in vivo, *Neuron* **104**, 655 (2019).
 - [6] W. L. Shew and D. Plenz, The functional benefits of criticality in the cortex, *Neuroscientist* **19**, 88 (2013).
 - [7] O. Kinouchi and M. Copelli, Optimal dynamical range of excitable networks at criticality, *Nat. Phys.* **2**, 348 (2006).
 - [8] W. L. Shew, H. Yang, T. Petermann, R. Roy, and D. Plenz, Neuronal avalanches imply maximum dynamic range in cortical networks at criticality, *J. Neurosci.* **29**, 15595 (2009).
 - [9] S. H. Gautam, T. T. Hoang, K. McClanahan, S. K. Grady, and W. L. Shew, Maximizing sensory dynamic range by tuning the cortical state to criticality, *PLoS Comput. Biol.* **11**, e1004576 (2015).
 - [10] W. L. Shew, H. Yang, S. Yu, R. Roy, and D. Plenz, Information capacity and transmission are maximized in balanced cortical networks with neuronal avalanches, *J. Neurosci.* **31**, 55 (2011).
 - [11] E. D. Fagerholm, G. Scott, W. L. Shew, C. Song, R. Leech, T. Knöpfel, and D. J. Sharp, Cortical entropy, mutual information and scale-free dynamics in waking mice, *Cerebral Cortex* **26**, 3945 (2016).
 - [12] W. P. Clawson, N. C. Wright, R. Wessel, and W. L. Shew, Adaptation towards scale-free dynamics improves cortical stimulus discrimination at the cost of reduced detection, *PLoS Comput. Biol.* **13**, e1005574 (2017).
 - [13] D. B. Larremore, W. L. Shew, E. Ott, F. Sorrentino, and J. G. Restrepo, Inhibition Causes Ceaseless Dynamics in Networks of Excitable Nodes, *Phys. Rev. Lett.* **112**, 138103 (2014).
 - [14] V. Agrawal, A. B. Cowley, Q. Alfaori, D. B. Larremore, J. G. Restrepo, and W. L. Shew, Robust entropy requires strong and balanced excitatory and inhibitory synapses, *Chaos* **28**, 103115 (2018).

- [31] A. Levina, J. M. Herrmann, and T. Geisel, Dynamical synapses causing self-organized criticality in neural networks, *Nat. Phys.* **3**, 857 (2007).
- [32] J. Hesse and T. Gross, Self-organized criticality as a fundamental property of neural systems, *Front. Syst. Neurosci.* **8**, 1 (2014).
- [33] Y. S. Virkar, J. G. Restrepo, W. L. Shew, and E. Ott, Dynamic regulation of resource transport induces criticality in interdependent networks of excitable units, *Phys. Rev. E* **101**, 022303 (2020).
- [34] R. Albert, H. Jeong, and A.-L. Barabási, Error and attack tolerance of complex networks, *Nature (London)* **406**, 378 (2000).