



## Ordering induced changes in the optical spectra of semiconductor alloys

J. E. Bernard, S. H. Wei, D. M. Wood, and Alex Zunger

Citation: [Applied Physics Letters](#) 52, 311 (1988); doi: 10.1063/1.99503

View online: <http://dx.doi.org/10.1063/1.99503>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/apl/52/4?ver=pdfcov>

Published by the [AIP Publishing](#)

---

Articles you may be interested in

[Structure and hyperfine parameters of cyclopropyl and bicyclobutyl radicals from post Hartree-Fock computations](#)

[J. Chem. Phys.](#) 104, 2630 (1996); 10.1063/1.470987

[Dependence of the optical properties of semiconductor alloys on the degree of long range order](#)

[Appl. Phys. Lett.](#) 62, 1937 (1993); 10.1063/1.109496

[Ordering in semiconductor alloys](#)

[Appl. Phys. Lett.](#) 56, 731 (1990); 10.1063/1.102695

[Band gap narrowing in ordered and disordered semiconductor alloys](#)

[Appl. Phys. Lett.](#) 56, 662 (1990); 10.1063/1.103307

[Elucidation of x ray diffraction data on the nature of the ordering of GaInP<sub>2</sub> and how the ordering relates to changes in the optical properties](#)

[Appl. Phys. Lett.](#) 54, 718 (1989); 10.1063/1.101470

---





