



7 ca dUf]gcb`cZ`h`Y`_`d`UbX`X]fYW`h`X]U[cbU`]nUh]cb`UddfcUW`Yg`hc`h`Y`Y`YW`fcb]Wghfi`Wi`fY`
cZ`b5g#`U5g`ei`Ubhi`a`Xchg

[L. W. Wang](#), [A. J. Williamson](#), [Alex Zunger](#), [H. Jiang](#), and [J. Singh](#)

Citation: [Applied Physics Letters](#) ⁺, 339 (io A9 (io



Comparison of the $\mathbf{k}\cdot\mathbf{p}$ and direct diagonalization approaches to the electronic structure of InAs/GaAs quantum dots

L. W. Wang, A. J. Williamson, and Alex Zunger^{a)}
National Renewable Energy Laboratory, Golden, Colorado 80401

H. Jiang and J. Singh
Department of Electrical Engineering and Computer Science, The University of Michigan, Ann Arbor, Michigan 48109-2122

Received 13 September 1999; accepted for publication 16 November 1999)

We present a comparison of the 8-band $\mathbf{k}\cdot\mathbf{p}$ and empirical pseudopotential approaches to describing the electronic structure of pyramidal InAs/GaAs self-assembled quantum dots. We find a generally good agreement between the two methods. The most significant shortcomings found in the $\mathbf{k}\cdot\mathbf{p}$ calculation are i) a reduced splitting of the electron p states (3 vs 24 meV), ii) an incorrect in-plane polarization ratio for electron-hole dipole transitions (0.97 vs 1.24), and iii) an over confinement of both electron (48 meV) and hole states (52 meV), resulting in a band gap error of 100 meV. We introduce a ‘‘linear combination of bulk bands’’ technique which produces results similar to a full direct diagonalization pseudopotential calculation, at a cost similar to the $\mathbf{k}\cdot\mathbf{p}$ method. © 2000 American Institute of Physics. S0003-6951(00)01903-3]

Self-assembled, Stranski–Krastanow (SK) grown semiconductor quantum dots such as InAs/GaAs have recently received considerable attention.¹ They exhibit a rich spec-

tionally expensive. In this approach, we do not limit the basis to $\psi_{n,k}$ -like states [Eq. 1)], but also include bulk Bloch functions, computed for a given value, ϵ , of the strain.

$$\psi_i^{\text{LCBB}}(\mathbf{r}) = \sum_n^{N_B} \sum_k^{N_k} C_{n,k}^{(i)} e^{i\mathbf{k}\cdot\mathbf{r}} u_{n,k}(\epsilon, \mathbf{r}), \quad 3)$$

where N_B and N_k are a cutoff for the number of bands and k points. The speed up of the LCBB method compared to the DD pseudopotential method arises from the fact that the LCBB states form a physically more intuitive basis than traditional plane waves and N_B and N_k can be significantly reduced to keep only the physically important bands and k points