

Finding the atomic configuration with a required physical property in multi-atom structures

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2007 J. Phys.: Condens. Matter 19 402201

(<http://iopscience.iop.org/0953-8984/19/40/402201>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 198.11.29.40

This content was downloaded on 04/03/2017 at 16:41

Please note that [terms and conditions apply](#).

You may also be interested in:

[Finding the lowest-energy crystal structure starting from randomly selected lattice vectors and atomic positions: first-principles evolutionary study of the Au–Pd, Cd–Pt, Al–Sc, Cu–Pd, Pd–Ti, and Ir–N binary systems](#)

Giancarlo Trimarchi and Alex Zunger

[A grand canonical genetic algorithm for the prediction of multi-component phase diagrams and testing of empirical potentials](#)

William W Tipton and Richard G Hennig

[Ab initio random structure searching](#)

Chris J Pickard and R J Needs

[Cluster expansions in multicomponent systems: precise expansions from noisy databases](#)

Alejandro Díaz-Ortiz, Helmut Dosch and Ralf Drautz

[Evolution of L12 ordered domains in fcc Cu₃Au alloy](#)

Mahdi Sanati and Alex Zunger

[Configurational thermodynamics of alloys from first principles: effective cluster interactions](#)

A V Ruban and I A Abrikosov



