




AD


D

$\Delta\mu = -3.2 \cdot 10^{-10}$ eV, $T = 14$ K (1200°), $p_2 = 1$ bar

Handwritten notes and calculations, including a table with columns labeled 1, 2, 3, 4 and rows of numerical data.

Handwritten notes and calculations, including a table with columns labeled 1, 2, 3, 4 and rows of numerical data.





(...), ...
 (...), ...
 2.5 ...
 3d ...
 13,41 ...
 2×10^4 ...
 0% ... 100

...
 2 | 4 ...
 4 ... 1.3 ...
 ... 120° ...
 41 2 ...
 ...
 10^{-3} ...
 10^{-1} ...
 ...
 ...
 ...
 ...

...
 4

$p_2 = 0.28$
 3×10^{20}
 10^{1-3}
 $p = 6 \times 10^{1-3}$
 $(1.3 \times 10^{1-3})^{11}$
 $(1.44 \times 10^{1-3})^{4,51}$
 $(0.203 \times 10^{1-3})^{52}$

10^{1-3}

4.5.2. Preliminary Synthesis of Li-Doped Cr_2MnO_4

1.1×10^{-2}
 $11,55$

13
 5
 13
 2
 4

This work was supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, Energy Frontier Research Centers, under Contract No. DE-AC36-08GO28308 to NREL. The high performance computing resources of the National Energy Research Scientific Computing Center and of NREL's Computational Science Center are gratefully acknowledged. This work made use of the J. B. Cohen X-Ray Diffraction Facility supported by the MRSEC program of the National Science Foundation (DMR-1121262) at the Materials Research Center of Northwestern University.

Received: March 5, 2013
 Published online: May 15, 2013

1.
